

Composition : Magolin-5 : Each film foated tablet contains Mirogabalin 5 mg as Mirogabalin Besylate INN.

Magolin-10 : Each film foated tablet contains Mirogabalin 10 mg as Mirogabalin Besylate INN.

Magolin -15 : Each film foated tablet contains Mirogabalin 15 mg as Mirogabalin Besylate INN.

Pharmacology : Mirogabalin has selective and potent binding affinities for human $\alpha\delta$ - subunits of VGCCs, which reduce calcium (Ca^{2+}) influx and neurotransmission in DRG, inhibiting neurotransmitter release in pre-synaptic neuron endings.

Indications : It is indicated to treatment of Neuropathic Pain, Diabetic peripheral neuropathic pain (DPNP), Postherpetic neuralgia (PHN), Peripheral neuropathic pain (PNP).

Dosage and administration : Normally, the initial dose for adults is 5 mg of Mirogabalin given orally twice daily, and then the dose is gradually increased by 5 mg at an interval of at least a week to 15 mg, given orally twice daily. The dose may be adjusted appropriately depending on patients ages and symptoms in the range of 10mg to 15mg or as directed by a registered physician.

Contraindication : Mirogabalin is contraindicated in patients with hypersensitive to Mirogabalin . Patients with mild to moderate hepatic & renal impairment.

Precaution : This medicine may cause dizziness, somnolence, or loss of consciousness. Avoid operating dangerous machinery, such as driving a car. Especially for elderly patients, careful attention should be taken. This medicine may cause weight gain. This medicine may cause blurred vision and double vision.

Side-effects : The most reported

Magolin

Mirogabalin Besylate
INN Tablet



**DRUG
INTERNATIONAL
LTD.**

adverse reactions include somnolence, dizziness, edema and weight gain. The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.

- Light headedness, state close to sleep with impaired consciousness, loss of consciousness (dizziness, somnolence, unconsciousness).
- General malaise, loss of appetite, nausea, vomiting, jaundice (liver dysfunction)

Use in Pregnancy and Lactation :

There is no information available on the safety profile on using in pregnancy & lactation .

Use in Child : There is no data available

Drug-Interactions : Mirogabalin does not induce or inhibit cytochrome P450 isoenzymes. Co administration of Mirogabalin with Cimetidine or Probenecid may raise the Mirogabalin plasma concentration. Importantly, if mirogabalin is taken with alcohol or lorazepam, the depressive effects on the CNS may be potentiated

Overdose : There is no data available.

Storage : Store at 30°C in a dry place.

Packing : Magolin -5 : Each box contains 14's tablets in a blister pack.

Magolin -10 : Each box contains 14's tablets in a blister pack.

Magolin -15 : Each box contains 14's tablets in a blister pack.