

# Fexofast

Fexofenadine Hydrochloride USP Tablet & Suspension

**Composition :**

**Fexofast - 60 :** Each film coated tablet contains Fexofenadine Hydrochloride USP 60mg.

**Fexofast -120 :** Each film coated tablet contains Fexofenadine Hydrochloride USP 120mg.

**Fexofast -180 :** Each film coated tablet contains Fexofenadine Hydrochloride USP 180mg.

**Fexofast Suspension :** Each 5ml suspension contains Fexofenadine Hydrochloride USP 30mg.

**Indications :** Fexofenadine Hydrochloride is indicated for the relief from physical symptoms associated with seasonal allergic rhinitis and treatment of chronic idiopathic urticaria. It is not cure totally but prevents the aggravation of rhinitis and urticaria. It reduces the severity of the symptoms and providing much relief from repeated sneezing, runny nose, itchy eyes and general body fatigue caused by rhinitis and urticaria.

**Dosage and administration : Fexofast tablets: Seasonal allergic rhinitis and chronic idiopathic urticaria:** Adults and Children 12 Years and Older: The recommended dose of Fexofenadine Hydrochloride is 60 mg twice daily or 180 mg once daily with water. A dose of 60 mg once daily is recommended as the starting dose in patients with decreased renal function. Children 6 to 11 Years: 30 mg twice daily with water. A dose of 30 mg once daily is recommended as the starting dose in patients with decreased renal function.

**Oral suspension: Seasonal allergic rhinitis:** Children 2 to 11 years: The recommended dose of oral suspension 30 mg twice daily. If pediatric patients decrease renal function, then adjust or maintain dose of 30 mg (5 ml) once daily recommended as the starting dose.

**Chronic idiopathic urticaria:** Children 6 months to 11 years: 30 mg (5 ml) twice daily for patients 2 to 11 years of age and 15 mg (2.5ml) twice daily for patients 6 months to less than 2 years of age. For pediatric patients with decrease renal function the recommended starting dose of oral suspension is 30 mg (5ml) once daily for patients 2 to 11 years of age and 15 mg (2.5ml) once daily for patients 6 months to less than 2 years of age. Shake bottle well, before each use. Or, as directed by the registered physician.

**Side effects :** Nausea, dizziness, vomiting, weakness, drowsiness, sleepiness, fatigue, diarrhea, unusual bowel movements, headache & fever.

**Contraindication :** Fexofenadine Hydrochloride is contraindicated in patients who are hypersensitive to this medication or its ingredients.

**Drug interactions :** Fexofenadine Hydrochloride should not be taken closely in time with aluminum and magnesium containing antacids. Co-administration of fexofenadine hydrochloride with either ketoconazole or erythromycin lead to increased plasma concentrations of fexofenadine in healthy adult subjects. Fexofenadine had no effect on the pharmacokinetics of either erythromycin or ketoconazole.

**Use in pregnancy and lactation :** There are no adequate and well-controlled studies in pregnant women. Fexofenadine Hydrochloride should be administered during pregnancy or lactation if clearly needed.

**Precautions :** Fexofenadine Hydrochloride should generally be avoided in patients with hepatic insufficiency. Patients with renal insufficiency should be given a lower initial dose.

**Overdose :** Dizziness, drowsiness, and dry mouth have been reported with fexofenadine hydrochloride overdose. Symptomatic and supportive treatment is recommended.

**Storage :** Store below 25° C in a dry place.

**Packing :**

**Fexofast - 60 :** Each box contains 5 x 10's tablets in blister pack.

**Fexofast -120 :** Each box contains 3 x 10's tablets in blister pack.

**Fexofast -180 :** Each box contains 2 x 10's tablets in blister pack.

**Fexofast Suspension :** Each bottle contains 50 ml Fexofenadine Hydrochloride USP suspension.

Manufactured by



**DRUG INTERNATIONAL LTD.**

Tongi, Gazipur, Bangladesh