

F-Son

Fluticasone Furoate INN 27.5 mcg/metered spray

Composition : F-Son Nasal Spray : Each metered spray contains Fluticasone Furoate INN 27.5 mcg.

Indications : This nasal spray indicated for the treatment of the symptoms of seasonal and perennial allergic rhinitis in patients aged 2 years and older.

Dosage and administration : Adults & Children over 12 years : 2 sprays in each nostril once a day. In some cases 2 sprays into each nostril twice daily, not exceeding 4 sprays.

Children under 12 years (2-11 Years) : 1 spray in each nostril once a day. Patients should use Fluticasone Furoate nasal spray at regular intervals as directed since its effectiveness depends on its regular use. Or, as directed by the registered physician.

Contraindication : Fluticasone Furoate nasal spray is contraindicated in patients with a hypersensitivity to any of its ingredients.

Precautions : Rarely, immediate hypersensitivity reactions or contact dermatitis may occur after the administration of Fluticasone Furoate nasal spray. Rare instances of wheezing, nasal septum perforation, cataracts, glaucoma and increased intraocular pressure have been reported following the intranasal application of corticosteroids, including Fluticasone Furoate. Although systemic effects have been minimal with recommended doses of Fluticasone Furoate nasal spray, potential risk increases with larger doses. Therefore, larger than recommended doses of Fluticasone Furoate nasal spray should be avoided.

Side effects : Side Effects are generally absorbed less into the rest of the body, therefore fewer side effects are seen. With the nasal spray, drying of the nose and an increase in the incidence of nosebleeds may occur.

Use in pregnancy and lactation : Pregnancy category C. There are no adequate and well controlled studies in pregnant women. Fluticasone Furoate nasal spray should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus. It is not known whether Fluticasone Furoate is excreted in human milk. However, other corticosteroids have been detected in human milk. Caution should be exercised when Fluticasone Furoate nasal spray is administered to a nursing woman.

Drug interactions : Potent inhibitors of (CYP3A4) may increase exposure to Fluticasone Furoate. Coadministration of ritonavir is not recommended. Use caution with coadministration of other potent CYP3A4 inhibitors, such as ketoconazole.

Overdose : There are no data on the effects of acute and chronic overdose with this Nasal Spray.

Storage : Store below 30°C, protect from light. Keep out of the reach of Children.

Packing : F-Son Nasal Spray : Each container contains 120 Metered doses for sprays.



Manufactured by

DRUG INTERNATIONAL LTD.

Tongi, Gazipur, Bangladesh